



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Viral infection causes sex-specific changes in fruit fly social aggregation behaviour**

**Citation for published version:**

Siva-Jothy, J & Vale, P 2019, 'Viral infection causes sex-specific changes in fruit fly social aggregation behaviour', *Biology letters*. <https://doi.org/10.1098/rsbl.2019.0344>

**Digital Object Identifier (DOI):**

[10.1098/rsbl.2019.0344](https://doi.org/10.1098/rsbl.2019.0344)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Biology letters

**Publisher Rights Statement:**

This is the accepted version of the following article: [full citation], which has been published in final form at [Link to final article].

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



1    **Viral infection causes sex-specific changes in fruit fly social aggregation behaviour**

2

3    Jonathon A. Siva-Jothy<sup>1</sup>, Pedro F. Vale<sup>1,2\*</sup>

4    <sup>1</sup>Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh,

5    EH9 3FL Edinburgh, United Kingdom

6    <sup>2</sup> Centre for Immunity, Infection and Evolution, University of Edinburgh, EH9 3FL Edinburgh,

7    United Kingdom

## 8    **Abstract**

9    Host behavioural changes following infection are common and could be important  
10    determinants of host behavioural competence to transmit pathogens. Identifying potential  
11    sources of variation in sickness behaviours is therefore central to our understanding of  
12    disease transmission. Here, we test how group social aggregation and individual locomotor  
13    activity vary between different genotypes of male and female fruit flies (*Drosophila*  
14    *melanogaster*) following septic infection with *Drosophila* C Virus (DCV). We find genetic-  
15    based variation in both locomotor activity and social aggregation, but we did not detect  
16    an effect of DCV infection on fly activity or sleep patterns within the initial days following  
17    infection. However, DCV infection caused sex-specific effects on social aggregation, as  
18    male flies in most genetic backgrounds increased the distance to their nearest neighbour  
19    when infected. We discuss possible causes for these differences in the context of individual  
20    variation in immunity and their potential consequences for disease transmission.

21

22    **Key words:** Social aggregation, locomotor activity, *Drosophila* Activity Monitor,  
23    *Drosophila*, *Drosophila* C Virus, sexual dimorphism

## 24 **Introduction**

25 Infection-induced changes to host physiology, and immunity in particular, following  
26 infection are well known, but it is equally striking that many animals experience similar  
27 behavioural changes following infection [1,2]. Common behavioural responses to infection  
28 include eating and moving less, as well as foregoing social and sexual interactions [1,3–5].  
29 Whether these behavioural changes in response to infection are evolved host responses,  
30 parasite manipulations, or a coincidental by-product of infection[6,7], they have potentially  
31 important consequences for disease transmission [8]. This is particularly clear for  
32 behaviours such as individual locomotor activity or group social aggregation, which will  
33 directly determine how frequently susceptible and infected individuals interact. Assessing  
34 how host behaviours that influence contact rates might change following infection is  
35 therefore central to understanding the spread of infectious disease.

36

37 The extent to which host behaviours are modified during infection is likely to depend on  
38 genetic and environmental factors. Even in the absence of infection, individuals of some  
39 genetic backgrounds are more likely to aggregate than others [9,10], while males and  
40 females in a broad range of host species often exhibit distinct behavioural profiles [11,12].  
41 How these different sources of variation influence infection-induced behavioural changes

42 is relatively understudied [8]. Measuring how males and females of different genetic  
43 backgrounds modify their behaviour during infection may highlight groups of individuals  
44 with higher contact rates and offer insight into the potential causes of heterogeneity in  
45 pathogen spread.

46

47 Testing if locomotor and aggregation behaviours change following infection, and if these  
48 changes differ between genetic backgrounds, is not straightforward for most host species.  
49 It requires knowledge of how individuals within a population differ in their genetic  
50 backgrounds and the ability to expose many individuals of the same background to  
51 infection in controlled conditions, while comparing their behavioural responses to infection  
52 with individuals of the same background that do not experience infection. For many animal  
53 species, and certainly in human populations, this type of intervention is either extremely  
54 challenging or not feasible. One alternative is to leverage the tools offered by model  
55 systems. *Drosophila melanogaster*, for example, has been widely used as a model system  
56 for behavioural genetics [13,14], and used specifically to study social aggregation and  
57 locomotor activity [9,15,16]. Further, *D. melanogaster* is a powerful model of immunity in  
58 response to a range of bacterial and viral pathogens [17]. Previous work has shown that  
59 *D. melanogaster* exhibits a range of behavioural changes following *Drosophila C Virus*

60 (DCV) infection, including pathogen avoidance during oviposition [18], foraging [19] and  
61 changes in locomotor [20–22]. Here, we test whether DCV infection changes social  
62 aggregation and locomotor activity in *D. melanogaster*, and if these effects vary with  
63 genetic background and sex.

## 64 **Materials & Methods**

### 65 **Flies and Rearing Conditions**

66 We used males and females from 10 lines sourced from the Drosophila Genetic Resource  
67 Panel (DGRP) [23], which are among the most and least susceptible genetic backgrounds  
68 to systemic Drosophila C Virus infection [24]. Genetic variation in DCV susceptibility was  
69 confirmed in a separate experiment where survival was measured following DCV infection  
70 in males and females from these lines (Figure S1; Table S3). Flies were reared in plastic  
71 vials on a standard diet of Lewis medium at  $18\pm1^{\circ}\text{C}$  with a 12 hour light:dark cycle with  
72 stocks tipped into new vials every 14 days. One month before the experiment, flies were  
73 transferred to incubators and maintained at  $25\pm1^{\circ}\text{C}$  with a 12 hour light:dark cycle at low  
74 density ( $\sim 10$  flies per vial) for two generations.

75

### 76 **Virus Culture and Infection**

77 The Drosophila C Virus (DCV) isolate was originally isolated in Charolles, France [25] and  
78 the stock used in this experiment was grown in Schneider Drosophila Line 2 (DL2) as  
79 previously described [20], diluted one hundred-fold ( $10^8$  infectious units per ml) in TRIS-  
80 HCl solution (pH=7.3), aliquoted and frozen at  $-70^{\circ}\text{C}$  until required. Given the extensive  
81 dilution of DL2 cells in TRIS buffer, 100% TRIS buffer was used as a control for the infection

solution. It is important to note that while our laboratory stocks are routinely screened for viruses and contaminants, unknown contaminants may be harboured by the DL2 cells. However, given the many orders of magnitude in the titers of possible contaminants compared to the titre of DCV, it is unlikely that these would cause effects confounded with the effect of DCV. To infect with DCV, 3-5-day old flies were pricked in the thorax in the mesopleura with a 0.15mm diameter pin, bent at 90° ~0.5mm from the tip, dipped in DCV (or TRIS-HCl for controls). Using this infection protocol establishes a systemic infection that results in increased viral titres within the first 3 days of infection [22,26–28].

## **Measuring *Drosophila* social aggregation**

Social aggregation was measured in a separate experiment, by calculating the nearest neighbour distance (NND) between individuals within a 12-fly group of the same sex and genetic background that were contained within a Petri dish for 30 minutes [10,16,29]. The experiment was conducted over five experimental blocks, each carried out over a single day, where each genetic background, sex and infection treatment was measured. Flies in infected treatment groups, were pricked with DCV 72 hours before their NND was measured. The NND was calculated by image analysis of pictures recorded of each group using the 'NND' package in ImageJ [30]. In total, we measured social aggregation in 580



groups of flies, with n=14-16 replicate groups of 12 flies for each genetic background, sex and infection status combination. To consider the effect of body size on social aggregation, we also measured the body length of a subset of individuals from each treatment group (Figure S1). NND measures were converted from millimetres to body lengths by dividing values by the average body length of individuals from treatment groups (Figure S2). A more detailed description of the experimental setup and analysis can be found in electronic supplementary material.

### **Measuring *Drosophila* activity**

The activity of single flies was measured during 4 continuous days using a *Drosophila* Activity Monitor (DAM2 System, TRIKinetics), in an incubator maintained at 25°C in a 12:12 light:dark cycle [15]. Over the course of the experiment, we measured the activity of 872 flies, with n=18-28 flies for each combination of sex and genetic background (Table S1). Raw activity data was processed using the DAM System File Scan Software [15], and the resulting data was manipulated using Microsoft Excel. We analysed fly activity data using three metrics: total locomotor activity, proportion of time spent asleep and the average activity when awake, as described previously [20]. A more detailed description of the experimental setup and analysis can be found in electronic supplementary material.

118

## 119 **Statistical Analysis**

120 We tested if differences in locomotor activity and social aggregation could be attributed  
121 to fly genetic background or sex using Generalized Linear Models (GLMs). Models used a  
122 full factorial 3-way interaction between infection status (control/infected), sex  
123 (male/female) and DGRP line (10 lines), all modelled as fixed effects. Analysis of social  
124 aggregation used a model listing only the median nearest neighbour distance of each dish  
125 as its response variable. To assess locomotor activity, we analysed 3 response variables in  
126 separate GLMs (total activity, proportion of time asleep, awake activity), adjusting the  
127 significance threshold to 0.01667 using Bonferroni correction to account for multiple-  
128 testing. All statistical analyses and graphics were carried out and produced in R 3.3.0 [31]  
129 using the *ggplot2* [32] and *lme4* [33] packages.

## Results

### Social aggregation

We found a significant effect of genetic background on the median nearest neighbour distances (NND) (Figure 1; Table 1). We found no evidence of sexual dimorphism in social aggregation across multiple genetic backgrounds, with no significant interaction between sex and genetic background. However, we observed that while female aggregation was not affected by infection, infected males aggregated further apart from each other compared to uninfected males (Figure 1; Table 1). This increase in the NND following infection was generally observed in males, regardless their genetic background (Figure 1; Table 1). We also detected an expected sexual dimorphism in body size, as female *D. melanogaster* are typically larger than males (Figure S1, Table S3). Incorporating this size difference into measures of social aggregation, by measuring body lengths between individuals did not alter the results qualitatively (Figure S2, Table S4).

### Locomotor activity

All three parameters of total locomotor activity, the proportion of time spent asleep and the average activity when awake, were affected by a combination of sex and genetic background (Figures 2 and S3; Table 2). However, there was no detectable difference in

148 how much infected and healthy flies moved or slept, and hence no evidence that infection  
149 impacted on any parameter of fly locomotor activity (Figures 2 and S3; Table 2).

150

151

## 152 **Discussion**

153 Identifying changes in host behaviour following infection is important to understand  
154 heterogeneity in disease transmission. Overall, our results indicate a significant sex  
155 difference in the effect of infection on social aggregation but no effect of infection on  
156 locomotor activity in either sex.

157

158 We observed that how closely flies aggregate with one another differs with their genetic  
159 background. The genetic variation we observed is similar to other studies that have  
160 measured nearest neighbour distance [10], as well as other aspects of *Drosophila* social  
161 behaviour, such as group size preference [9] and group composition [34]. Group  
162 composition is affected by the natural *foraging* gene polymorphism, where larvae are either  
163 sitters, which aggregate toward conspecifics at food sources or rovers, who are more prone  
164 to independent food searching behaviour. Larger groups of larvae on food patches are  
165 more likely to be comprised of sitters, as rovers leave food patches after overcrowding

[34]. Genetic components of social behaviour have also been identified in a number of mammal species, including humans [35]. In a number of vole species, variation in oxytocin [36] and arginine vasopressin [37] receptor density is associated with between-species variation in pair-bonding and monogamy.

While aggregation between healthy males and females did not differ, once infected, males moved further apart from one another, while female aggregation did not change. One possible explanation for why males aggregate further apart following infection is a sex difference in immunity and the costs of social aggregation [38]. Sexually dimorphic immunity may be particularly relevant as male *D. melanogaster* exhibit a stereotyped suite of aggressive behaviours [39–41]. While fighting can gain males access to valuable resources, it often incurs substantial costs [42,43]. DCV infection could exacerbate the cost of male aggregation, as resources would also need to be spent on fighting infection, which could lead to males aggregating less. Despite females also fighting one another, this aggression is generally less costly [44,45]. Females may therefore still be able to aggregate relatively closely while fighting DCV infection.

183 Irrespective of the metric used, we found no measurable effect of DCV infection on  
184 locomotor activity. Other work has shown decreases in *Drosophila* daily movement  
185 following injection with DCV, where a marked reduction is seen after 4 days of infection  
186 [21]. Reduced daily locomotor activity has also been observed in *Drosophila* after 3 days  
187 of infection with the DNA virus Kalithea virus [46]. Injecting, rather than pricking, flies with  
188 viral suspension, allows more precise control of infectious dosage, which could also  
189 increase infection severity [47]. Another potential explanation is that we infected flies via  
190 thoracic pricking, as opposed to abdominal injection which has been shown to reduce  
191 resistance to infection in *Drosophila* [48]. The injury produced by thoracic pricking may  
192 obscure subtler changes to activity produced by DCV infection. Orally infecting flies shows  
193 a range of sex-specific behavioural symptoms, with sub-lethal doses reducing daily  
194 locomotor activity in males after 3-6 days of infection [22]. Conversely, following oral  
195 infection with larger doses of DCV, females, but not males, have been shown to sleep  
196 more [20]. These studies suggest we may not have seen an effect of DCV infection on  
197 activity, because infections were not severe enough to elicit behavioural symptoms.  
198 Measuring the activity of flies later in infection might address these explanations, as this  
199 will enable flies to heal from thoracic injury and accrue a greater viral burden.

200

201 We measured social aggregation in groups of individuals composed of the same genetic  
202 background, sex and infection status in order to dissect their influence on social  
203 aggregation. However, in more heterogenous wild populations these characteristics can  
204 produce population structure that could affect contact between individuals. Individuals  
205 with shared genotypes can be more likely to interact due to predispositions to traits such  
206 as group size preference [34,49] and aggression [50]. Similarly, sexual interactions between  
207 males and females, as well as fighting and other forms of sexual competition, further alter  
208 contact networks within populations [51,52]. When present together, healthy hosts might  
209 also be able to avoid infected conspecifics by detecting the pathogen or cues of its  
210 pathology [53]. Future work aiming to characterise the influence of these sources of  
211 variation on heterogeneity in contact rate should consider how they change with, and are  
212 changed by, population structure.

213

214 The contrasting ways social aggregation and locomotor activity change following infection  
215 highlight the complexity of sources determining between-individual variation in disease  
216 transmission. This is complicated further by sex differences across and within these genetic  
217 backgrounds. The change induced by DCV infection on social aggregation but not  
218 locomotor activity also demonstrates the importance of considering multiple host

219 behaviours. Central to understanding the effect of this genetic and sex-specific variation  
220 in social aggregation and locomotor activity on heterogeneity in disease transmission is  
221 characterising their effect on contact rates. Additionally, future work should consider how  
222 these traits interact with other key determinants of transmission, such as infectiousness  
223 and infection duration, as these three components ultimately define disease transmission  
224 in conjunction with one another.

225

## 226 **Acknowledgements**

227 We are grateful to Katy Monteith for general lab assistance, Angela Reid and Lucinda  
228 Rowe for preparation of all fly media and the Regan, Obbard, Walling and Vale lab  
229 members for constructive discussion of these ideas.

230

## 231 **Funding**

232 J.A.S.J. was supported by a NERC E3 DTP PhD studentship. P.F.V. was supported by a  
233 Branco Weiss fellowship (<https://brancoweissfellowship.org/>) and a Chancellor's Fellowship  
234 (School of Biological Sciences, University of Edinburgh).

235

236 Data Accessibility:



237 All data and R code can be accessed on Data Dryad:

238 <https://datadryad.org/review?doi=doi:10.5061/dryad.9232648>

239 DOI: <https://doi.org/10.5061/dryad.9232648>

240

241

242 Authors' contributions:

243 JSJ and PFV conceived and designed the study; JSJ carried out the experimental work,

244 acquired and analysed the and drafted the manuscript; JSJ and PFV wrote the

245 manuscript. JSJ approved the final version to be published. JSJ and PFV agree to be

246 accountable for all aspects of the work in ensuring that questions related to the accuracy

247 or integrity of any part of the work are appropriately investigated and resolved.

248

249

250

251

252 **Figure 1** – Mean $\pm$ SE median nearest neighbour distance (NND) in millimetres (mm). (a)  
253 Uninfected female-only arenas shown in blue, and infected female-only bars in pale blue.  
254 (b) Uninfected male-only arenas are shown in red, and infected male-only arenas in pink.  
255 The x-axis of both panels is ordered from the lowest to highest mean median NND of  
256 female flies.

257

258

259 **Figure 2** – Mean $\pm$ SE (A) total locomotor activity, (B) proportion of time flies spent  
260 sleeping and (C) mean activity while flies were awake, during the first 96 hours following  
261 DCV infection. Across all panels, sex and infection status are represented by colour with  
262 uninfected females shown in blue, infected females in pale blue, uninfected males in red,  
263 and infected males in pink. The order of genetic backgrounds on the x-axis of each of  
264 panel follows the ascending order of female flies.

265 **Tables**

| Response   | Predictor                               | Df | F       | p          |
|------------|---|----|---------|------------|
| Median NND | Genetic Background                      | 9  | 5.0249  | <0.0001*** |
|            | Sex                                     | 1  | 2.7870  | 0.13       |
|            | Infection                               | 1  | 21.1301 | <0.0001*** |
|            | Genetic Background<br>× Sex             | 9  | 1.4112  | 0.19       |
|            | Genetic Background<br>× Infection       | 9  | 0.9654  | 0.49       |
|            | Sex × Infection                         | 1  | 19.6600 | <0.0001*** |
|            | Genetic Background<br>× Sex × Infection | 9  | 1.6729  | 0.12       |

266

267 Table 1 - Model outputs for statistical tests performed on social aggregation, testing the  
 268 causes of variation in sociality in males and females of 10 *D. melanogaster* genetic  
 269 backgrounds. Significant predictors are marked with asterisks (p<0.05=\*, p<0.01=\*\* and  
 270 p<0.001=\*\*\*).

271

| Response                        | Predictor                            | Df | F       | p          |
|---------------------------------|--------------------------------------|----|---------|------------|
| Total Activity                  | Genetic Background                   | 9  | 14.83   | <0.0001*** |
|                                 | Sex                                  | 1  | 1.537   | 0.21       |
|                                 | Infection                            | 1  | 0.117   | 0.73       |
|                                 | Genetic Background × Sex             | 9  | 3.0485  | 0.0013*    |
|                                 | Genetic Background × Infection       | 9  | 1.4125  | 0.18       |
|                                 | Sex × Infection                      | 1  | 3.9707  | 0.047      |
|                                 | Genetic Background × Sex × Infection | 9  | 1.9471  | 0.043      |
| Proportion of Time Spent Asleep | Genetic Background                   | 9  | 25.1759 | <0.0001*** |
|                                 | Sex                                  | 1  | 77.9823 | <0.0001*** |
|                                 | Infection                            | 1  | 0.6939  | 0.41       |
|                                 | Genetic Background × Sex             | 9  | 3.444   | <0.001**   |
|                                 | Genetic Background × Infection       | 9  | 0.8021  | 0.61       |
|                                 | Sex × Infection                      | 1  | 0.7513  | 0.39       |
|                                 | Genetic Background × Sex × Infection | 9  | 1.4612  | 0.16       |
| Awake Activity                  | Genetic Background                   | 9  | 8.1673  | <0.0001*** |
|                                 | Sex                                  | 1  | 0.6641  | 0.54       |
|                                 | Infection                            | 1  | 0.0008  | 0.86       |
|                                 | Genetic Background × Sex             | 9  | 5.2153  | 0.0013*    |
|                                 | Genetic Background × Infection       | 9  | 0.8716  | 0.58       |
|                                 | Sex × Infection                      | 1  | 0.8430  | 0.44       |
|                                 | Genetic Background × Sex × Infection | 9  | 1.2998  | 0.61       |

272

273 **Table 2** – Model outputs for statistical tests performed on host activity data, testing the  
274 causes of variation in locomotor activity, sleep patterns and average awake activity in males  
275 and females of 10 *D. melanogaster* genetic backgrounds. Significance thresholds are  
276 corrected for multiple testing using Bonferroni correction, with significant predictors are  
277 marked with asterisks ( $p < 0.01667 = *$ ,  $p < 0.001 = **$  and  $p < 0.0001 = ***$ ).

## References

1. Hart BL. 1988 Biological basis of the behavior of sick animals. *Neuroscience & Biobehavioral Reviews* **12**, 123–137. (doi:10.1016/S0149-7634(88)80004-6)
2. Poulin R. 1995 ‘Adaptive’ changes in the behaviour of parasitized animals: a critical review. *Int. J. Parasitol.* **25**, 1371–1383.
3. Kazlauskas N, Klappenbach M, Depino AM, Locatelli FF. 2016 Sickness Behavior in Honey Bees. *Front Physiol* **7**. (doi:10.3389/fphys.2016.00261)
4. Hart BL. 2011 Behavioural defences in animals against pathogens and parasites: parallels with the pillars of medicine in humans. *Phil. Trans. R. Soc. B* **366**, 3406–3417. (doi:10.1098/rstb.2011.0092)
5. Lopes PC. 2014 When is it socially acceptable to feel sick? *Proceedings of the Royal Society of London B: Biological Sciences* **281**, 20140218. (doi:10.1098/rspb.2014.0218)
6. de Roode JC, Lefèvre T. 2012 Behavioral Immunity in Insects. *Insects* **3**, 789–820. (doi:10.3390/insects3030789)
7. Vale PF, Siva-Jothy J, Morrill A, Forbes M. 2018 The influence of parasites on insect behavior. In *Insect Behavior: from mechanisms to ecological and evolutionary consequences*, pp. 274–291. OUP Oxford.
8. Barron DG, Gervasi SS, Pruitt JN, Martin LB. 2015 Behavioral competence: How host behaviors can interact to influence parasite transmission risk. *Current Opinion in Behavioral Sciences* **6**, 35–40. (doi:10.1016/j.cobeha.2015.08.002)
9. Saltz JB. 2011 NATURAL GENETIC VARIATION IN SOCIAL ENVIRONMENT CHOICE: CONTEXT-DEPENDENT GENE-ENVIRONMENT CORRELATION IN DROSOPHILA MELANOGASTER. *Evolution* **65**, 2325–2334. (doi:10.1111/j.1558-5646.2011.01295.x)
10. Anderson BB, Scott A, Dukas R. 2016 Social behavior and activity are decoupled in larval and adult fruit flies. *Behavioral Ecology* **27**, 820–828. (doi:10.1093/beheco/arv225)
11. Manoli DS, Foss M, Villella A, Taylor BJ, Hall JC, Baker BS. 2005 Male-specific *fruitless* specifies the neural substrates of *Drosophila* courtship behaviour. *Nature* **436**, 395–400. (doi:10.1038/nature03859)
12. Walum H *et al.* 2008 Genetic variation in the vasopressin receptor 1a gene (AVPR1A) associates with pair-bonding behavior in humans. *PNAS* **105**, 14153–14156. (doi:10.1073/pnas.0803081105)
13. Dubnau J. 2014 *Behavioral Genetics of the Fly (Drosophila Melanogaster)*. Cambridge University Press.
14. Sokolowski MB. 2001 *Drosophila*: genetics meets behaviour. *Nature reviews. Genetics* **2**, 879–90. (doi:10.1038/35098592)

- 315 15. Pfeifferberger C, Lear BC, Keegan KP, Allada R. 2010 Processing Sleep Data Created  
316 with the Drosophila Activity Monitoring (DAM) System. *Cold Spring Harb Protoc* **2010**,  
317 pdb.prot5520. (doi:10.1101/pdb.prot5520)
- 318 16. Simon AF, Chou M-T, Salazar ED, Nicholson T, Saini N, Metchev S, Krantz DE. 2012 A  
319 simple assay to study social behavior in Drosophila: measurement of social space within  
320 a group. *Genes, brain, and behavior* **11**, 243–52. (doi:10.1111/j.1601-  
321 183X.2011.00740.x)
- 322 17. Troha K, Buchon N. 2019 Methods for the study of innate immunity in Drosophila  
323 melanogaster. *Wiley Interdiscip Rev Dev Biol* , e344. (doi:10.1002/wdev.344)
- 324 18. Siva-Jothy JA, Monteith KM, Vale PF. 2018 Navigating infection risk during oviposition  
325 and cannibalistic foraging in a holometabolous insect. *Behav Ecol*  
326 (doi:10.1093/beheco/ary106)
- 327 19. Vale PF, Jardine MD. 2016 Viral exposure reduces the motivation to forage in female  
328 Drosophila melanogaster. *Fly* , 1–7. (doi:10.1080/19336934.2016.1207029)
- 329 20. Vale PF, Jardine MD. 2015 Sex-specific behavioural symptoms of viral gut infection and  
330 Wolbachia in Drosophila melanogaster. *Journal of Insect Physiology* **82**, 28–32.  
331 (doi:10.1016/J.JINSPHYS.2015.08.005)
- 332 21. Arnold PA, Johnson KN, White CR. 2013 Physiological and metabolic consequences of  
333 viral infection in Drosophila melanogaster. *The Journal of Experimental Biology* **216**,  
334 3350–3357. (doi:10.1242/jeb.088138)
- 335 22. Gupta V, Stewart CO, Rund SSC, Monteith K, Vale PF. 2017 Costs and benefits of  
336 sublethal Drosophila C virus infection. *Journal of Evolutionary Biology* **30**, 1325–1335.  
337 (doi:10.1111/jeb.13096)
- 338 23. Mackay TFC *et al.* 2012 The Drosophila melanogaster Genetic Reference Panel. *Nature*  
339 **482**, 173–8. (doi:10.1038/nature10811)
- 340 24. Magwire MM, Fabian DK, Schweyen H, Cao C, Longdon B, Bayer F, Jiggins FM. 2012  
341 Genome-Wide Association Studies Reveal a Simple Genetic Basis of Resistance to  
342 Naturally Coevolving Viruses in Drosophila melanogaster. *PLoS Genetics* **8**.  
343 (doi:10.1371/journal.pgen.1003057)
- 344 25. Jousset F-X, Bergoin M, Revet B. 1977 Characterization of the Drosophila C Virus.  
345 *Journal of General Virology* **34**, 269–283. (doi:10.1099/0022-1317-34-2-269)
- 346 26. Gupta V, Vale PF. 2017 Nonlinear disease tolerance curves reveal distinct components of  
347 host responses to viral infection. *R Soc Open Sci* **4**, 170342. (doi:10.1098/rsos.170342)
- 348 27. Dostert C, Jouanguy E, Irving P, Troxler L, Galiana-Arnoux D, Hetru C, Hoffmann JA,  
349 Imler J-L. 2005 The Jak-STAT signaling pathway is required but not sufficient for the  
350 antiviral response of drosophila. *Nat Immunol* **6**, 946–953. (doi:10.1038/ni1237)
- 351 28. Rij RP van, Merklings SH. 2015 Analysis of resistance and tolerance to virus infection in  
352 Drosophila. *Nature Protocols* **10**, 1084. (doi:10.1038/nprot.2015.071)

- 353 29. Clark PJ, Evans FC. 1954 Distance to Nearest Neighbor as a Measure of Spatial  
354 Relationships in Populations. *Ecology* **35**, 445–453. (doi:10.2307/1931034)
- 355 30. Schneider CA, Rasband WS, Eliceiri KW. 2012 NIH Image to ImageJ: 25 years of Image  
356 Analysis. *Nat Methods* **9**, 671–675.
- 357 31. R Core Team. 2017 R: A language and environment for statistical computing. *R*  
358 *Foundation for Statistical Computing, Vienna, Austria*
- 359 32. Wickham H. 2016 *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New  
360 York.
- 361 33. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models  
362 Using **lme4**. *Journal of Statistical Software* **67**. (doi:10.18637/jss.v067.i01)
- 363 34. Philippe A-S, Jeanson R, Pasquaretta C, Rebaudo F, Sueur C, Mery F. 2016 Genetic  
364 variation in aggregation behaviour and interacting phenotypes in *Drosophila*.  
365 *Proceedings. Biological sciences* **283**, 20152967–20152967.  
366 (doi:10.1098/rspb.2015.2967)
- 367 35. Anacker AMJ, Beery AK. 2013 Life in groups: the roles of oxytocin in mammalian  
368 sociality. *Front Behav Neurosci* **7**. (doi:10.3389/fnbeh.2013.00185)
- 369 36. Ophir AG, Gessel A, Zheng D-J, Phelps SM. 2012 Oxytocin receptor density is  
370 associated with male mating tactics and social monogamy. *Hormones and Behavior* **61**,  
371 445–453. (doi:10.1016/j.yhbeh.2012.01.007)
- 372 37. Winslow JT, Hastings N, Carter CS, Harbaugh CR, Insel TR. 1993 A role for central  
373 vasopressin in pair bonding in monogamous prairie voles. *Nature* **365**, 545.  
374 (doi:10.1038/365545a0)
- 375 38. Kelly CD, Stoehr AM, Nunn C, Smyth KN, Prokop ZM. 2018 Sexual dimorphism in  
376 immunity across animals: a meta-analysis. *Ecol. Lett.* **21**, 1885–1894.  
377 (doi:10.1111/ele.13164)
- 378 39. Dierick H a, Greenspan RJ. 2006 Molecular analysis of flies selected for aggressive  
379 behavior. *Nature genetics* **38**, 1023–31. (doi:10.1038/ng1864)
- 380 40. Dierick HA. 2007 A method for quantifying aggression in male *Drosophila melanogaster*.  
381 *Nature protocols* **2**, 2712–8. (doi:10.1038/nprot.2007.404)
- 382 41. Baxter CM, Barnett R, Dukas R. 2015 Aggression, mate guarding and fitness in male  
383 fruit flies. *Animal Behaviour* **109**, 235–241. (doi:10.1016/j.anbehav.2015.08.023)
- 384 42. Chen S, Lee AY, Bowens NM, Huber R, Kravitz EA. 2002 Fighting fruit flies : A model  
385 system for the study of aggression. *Proceedings of the National Academy of Sciences of*  
386 *the United States of America* **99**, 5664–5668.
- 387 43. Lim RS, Eyj?lfsd??ttir E, Shin E, Perona P, Anderson DJ. 2014 How food controls  
388 aggression in *Drosophila*. *PLoS ONE* **9**. (doi:10.1371/journal.pone.0105626)



44. Ueda A, Kidokoro Y. 2002 Aggressive behaviours of female *Drosophila melanogaster* are influenced by their social experience and food resources. *Physiological Entomology* **27**, 21–28. (doi:10.1046/j.1365-3032.2002.00262.x)
45. Nilsen SP, Chan Y-B, Huber R, Kravitz EA. 2004 Gender-selective patterns of aggressive behavior in *Drosophila melanogaster*. *PNAS* **101**, 12342–12347. (doi:10.1073/pnas.0404693101)
46. Palmer WH, Medd NC, Beard PM, Obbard DJ. 2018 Isolation of a natural DNA virus of *Drosophila melanogaster*, and characterisation of host resistance and immune responses. *PLOS Pathogens* **14**, e1007050. (doi:10.1371/journal.ppat.1007050)
47. Neyen C, Bretscher AJ, Binggeli O, Lemaitre B. 2014 Methods to study *Drosophila* immunity. *Methods (San Diego, Calif.)* **68**, 116–28. (doi:10.1016/j.ymeth.2014.02.023)
48. Chambers MC, Jacobson E, Khalil S, Lazzaro BP. 2014 Thorax Injury Lowers Resistance to Infection in *Drosophila melanogaster*. *Infect Immun* **82**, 4380–4389. (doi:10.1128/IAI.02415-14)
49. Brown CR, Brown MB. 2000 Heritable basis for choice of group size in a colonial bird. *PNAS* **97**, 14825–14830. (doi:10.1073/pnas.97.26.14825)
50. Lea AJ, Blumstein DT, Wey TW, Martin JGA. 2010 Heritable victimization and the benefits of agonistic relationships. *Proc Natl Acad Sci U S A* **107**, 21587–21592. (doi:10.1073/pnas.1009882107)
51. Perkins SE, Ferrari MF, Hudson PJ. 2008 The effects of social structure and sex-biased transmission on macroparasite infection. *Parasitology* **135**, 1561–1569. (doi:10.1017/S0031182008000449)
52. Silk MJ, Weber NL, Steward LC, Hodgson DJ, Boots M, Croft DP, Delahay RJ, McDonald RA. 2018 Contact networks structured by sex underpin sex-specific epidemiology of infection. *Ecology Letters* **21**, 309–318. (doi:10.1111/ele.12898)
53. Curtis VA. 2014 Infection-avoidance behaviour in humans and other animals. *Trends in Immunology* **35**, 457–464. (doi:10.1016/j.it.2014.08.006)